

Report of the College of Physicians for Assisted Reproduction Therapy

Belgium 2008

16 September 2010

Version 1.0

Table of Contents

Section 1: General overview	6
Table 1.1 All cycles: Type of cycles.....	6
Table 1.2 Own and recipient fresh cycles: Number of laboratory treatment cycles	7
Figure 1.3 Own and recipient fresh cycles: Evolution of total number of cycles	8
Section 2: own fresh cycles.....	9
Table 2.1 Own fresh cycles: Overview of cycles	9
Figure 2.2 Own fresh cycles: Female age and laborank.....	10
Figure 2.3 Own fresh cycles: Indications of ART.....	11
Table 2.4 Own fresh cycles: Indications of ART: female and male causes	12
Figure 2.5 Own fresh cycles: Female age distribution	13
Figure 2.6 Own fresh cycles: Pituitary inhibition	14
Table 2.7 Own fresh cycles: Stimulation protocol	15
Figure 2.8 Own fresh cycles: Total dose of Gonadotrophins (percentiles).....	16
Figure 2.9 Own fresh cycles: Total dose of Gonadotrophins (boxplot)	17
Table 2.10 Own fresh cycles: Methods of fertilization	18
Table 2.11 Own fresh cycles: ICSI method sperm from partner.....	19
Table 2.12 Own fresh cycles: Transfers by age and rank categories	20
Table 2.13 Own fresh cycles: Transfers by social security	21
Figure 2.14 Own fresh cycles: Embryos transferred women < 36 years old	22
Figure 2.15 Own fresh cycles: Embryos transferred women 36-40 years old	23
Figure 2.16 Own fresh cycles: Embryos transferred women 40-43 years old	24
Table 2.17 Own fresh cycles: Laboratory data	25
Figure 2.18 Own fresh cycles: Summary pick-up cycles	26
Figure 2.19 Own fresh cycles: Distribution of embryo transfers	27
Table 2.20 Own fresh cycles: Cause of no transfer	28
Figure 2.21 Own fresh cycles: Day of embryos transfer	29
Table 2.22 Own fresh cycles: Cycles with cryopreservation	30
Table 2.23 Own fresh cycles: Number of HCG+ pregnancies according to age and rank	31
Table 2.24 Own fresh cycles: Number of clinical pregnancies according to age and rank.....	35
Table 2.25 Own fresh cycles: Number of clinical pregnancies including FHB according to age and rank	39
Table 2.26 Own fresh cycles: Number of deliveries according to age and rank.....	43
Figure 2.27 Own fresh cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age	47
Figure 2.28 Own fresh cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age.....	48
Figure 2.29 Own fresh cycles: Birth rate per transferred embryo according to age	49
Figure 2.30 Own fresh cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to rank	50
Figure 2.31 Own fresh cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to rank	51
Figure 2.32 Own fresh cycles: Birth rate per transferred embryo according to rank.....	52
Figure 2.33 Own fresh cycles: Number of deliveries	53

Figure 2.34 Own fresh cycles: Type of deliveries	54
Table 2.35 Own fresh cycles: Sex of babies	55
Table 2.36 Own fresh cycles: Birth weight.....	56
Table 2.37 Own fresh cycles: Gestational age at delivery	57
Figure 2.38 Own fresh cycles: Birth weight (boxplot).....	58
Figure 2.39 Own fresh cycles: Gestational age at delivery (boxplot)	59
Table 2.40 Own fresh cycles: Prevalence of preterm birth according to type of delivery	60
Table 2.41 Own fresh cycles: Prevalence of low birth weight according to type of delivery	61
Figure 2.42 Own fresh cycles: Evolution of number of embryos transferred.....	62
Figure 2.43 Own fresh cycles: Evolution of number of single and multiple deliveries	63
Section 3: Own cryo cycles	64
Table 3.1 Own cryo cycles: Overview of cryo cycles.....	64
Table 3.2 Own cryo cycles: Number of embryos transferred	65
Table 3.3 Own cryo cycles: Pituitary inhibition	66
Table 3.4 Own cryo cycles: Stimulation protocol	67
Table 3.5 Own cryo cycles: Number of HCG+ pregnancies according to age.....	68
Table 3.6 Own cryo cycles: Number of clinical pregnancies according to age	69
Table 3.7 Own cryo cycles: Number of clinical pregnancies including FHB according to age	70
Table 3.8 Own cryo cycles: Number of deliveries according to age.....	71
Figure 3.9 Own cryo cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age	72
Figure 3.10 Own cryo cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age	73
Figure 3.11 Own cryo cycles: Birth rate per transferred embryo according to age.....	74
Figure 3.12 Own cryo cycles: Number of deliveries.....	75
Figure 3.13 Own cryo cycles: Type of deliveries.....	76
Table 3.14 Own cryo cycles: Sex of babies	77
Table 3.15 Own cryo cycles: Birth weight.....	78
Table 3.16 Own cryo cycles: Gestational age at delivery.....	79
Figure 3.17 Own cryo cycles: Birth weight (boxplot).....	80
Figure 3.18 Own cryo cycles: Gestational age at delivery (boxplot).....	81
Table 3.19 Own cryo cycles: Prevalence of preterm birth according to type of delivery	82
Table 3.20 Own cryo cycles: Prevalence of low birth weight according to type of delivery.....	83
Section 4: Fresh recipient cycles	84
Table 4.1 Fresh recipient cycles: Overview of cycles.....	84
Figure 4.2 Fresh recipient cycles: Female age and laborank	85
Figure 4.3 Fresh recipient cycles: Female age distribution	86
Figure 4.4 Fresh recipient cycles: Pituitary inhibition	87
Table 4.5 Fresh recipient cycles: Stimulation protocol.....	88
Table 4.6 Fresh recipient cycles: Number of HCG+ pregnancies according to age	89
Table 4.7 Fresh recipient cycles: Number of clinical pregnancies according to age.....	90
Table 4.8 Fresh recipient cycles: Number of clinical pregnancies including FHB according to age.....	91
Table 4.9 Fresh recipient cycles: Number of deliveries according to age.....	92

Figure 4.10 Fresh recipient cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age.....	93
Figure 4.11 Fresh recipient cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age.....	94
Figure 4.12 Fresh recipient cycles: Birth rate per transferred embryo according to age	95
Figure 4.13 Fresh recipient cycles: Number of deliveries	96
Table 4.14 Fresh recipient cycles: Type of deliveries.....	97
Table 4.15 Fresh recipient cycles: Sex of babies.....	98
Table 4.16 Fresh recipient cycles: Birth weight.....	99
Table 4.17 Fresh recipient cycles: Gestational age at delivery	100
Figure 4.18 Fresh recipient cycles: Birth weight (boxplot).....	101
Figure 4.19 Fresh recipient cycles: Gestational age at delivery (boxplot)	102
Table 4.20 Fresh recipient cycles: Prevalence of preterm birth according to type of delivery.....	103
Table 4.21 Fresh recipient cycles: Prevalence of low birth weight according to type of delivery	104
Section 5: Cryo recipient cycles.....	105
Table 5.1 Cryo recipient cycles: Overview of cryo cycles	105
Table 5.2 Cryo recipient cycles: Number of embryos transferred.....	106
Table 5.3 Cryo recipient cycles: Pituitary inhibition.....	107
Table 5.4 Cryo recipient cycles: Stimulation protocol.....	108
Table 5.5 Cryo recipient cycles: Number of HCG+ pregnancies according to age	109
Table 5.6 Cryo recipient cycles: Number of clinical pregnancies according to age.....	110
Table 5.7 Cryo recipient cycles: Number of clinical pregnancies including FHB according to age.....	111
Table 5.8 Cryo recipient cycles: Number of deliveries according to age	112
Figure 5.9 Cryo recipient cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age	113
Figure 5.10 Cryo recipient cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age	114
Figure 5.11 Cryo recipient cycles: Birth rate per transferred embryo according to age	115
Figure 5.12 Cryo recipient cycles: Number of deliveries	116
Table 5.13 Cryo recipient cycles: Type of deliveries.....	117
Table 5.14 Cryo recipient cycles: Sex of babies.....	118
Table 5.15 Cryo recipient cycles: Birth weight	119
Table 5.16 Cryo recipient cycles: Gestational age at delivery	120
Figure 5.17 Cryo recipient cycles: Birth weight (boxplot)	121
Figure 5.18 Cryo recipient cycles: Gestational age at delivery (boxplot).....	122
Table 5.19 Cryo recipient cycles: Prevalence of preterm birth according to type of delivery	123
Table 5.20 Cryo recipient cycles: Prevalence of low birth weight according to type of delivery	124
Section 6: Fresh donor cycles	125
Table 6.1 Fresh donor cycles: Overview of cycles	125
Figure 6.2 Fresh donor cycles: Female age distribution.....	126
Table 6.3 Fresh donor cycles: Pituitary inhibition	127
Table 6.4 Fresh donor cycles: Stimulation protocol	128
Figure 6.5 Fresh donor cycles: Total dose of Gonadotrophins (percentiles).....	129
Section 7: Appendix.....	130
Table 7.1 : Definitions.....	130

Table 7.2 : List of B-centres having supplied data131
Colophon132

Section 1: General overview

Table 1.1 All cycles: Type of cycles


	Statistic	Total (N=29541)	All Centres	
			With social security (N=24961)	Without social security (N=4580)
Own fresh cycle	n (%)	19061 (64.52%)	16211 (64.95%)	2850 (62.23%)
Own cryo cycle	n (%)	8576 (29.03%)	7610 (30.49%)	966 (21.09%)
Fresh recipient cycle	n (%)	820 (2.78%)	459 (1.84%)	361 (7.88%)
Cryo recipient cycle	n (%)	294 (1.00%)	157 (0.63%)	137 (2.99%)
Fresh Donor cycle	n (%)	578 (1.96%)	394 (1.58%)	184 (4.02%)
Fresh sharing cycle	n (%)	82 (0.28%)	32 (0.13%)	50 (1.09%)
Mixed (fresh + thawed) cycle	n (%)	24 (0.08%)	16 (0.06%)	8 (0.17%)
Unspecified fresh cycle	n (%)	12 (0.04%)	12 (0.05%)	0 (0.00%)
Unspecified cryo cycle	n (%)	93 (0.31%)	69 (0.28%)	24 (0.52%)
Unknown cycle type	n (%)	1 (0.00%)	1 (0.00%)	0 (0.00%)

Table 1.2 Own and recipient fresh cycles: Number of laboratory treatment cycles

	All Centres (N=16999, Missing=1078)		
	Patients with Social Security	Patients without Social Security	Total
	N (%)	N (%)	N
All ages & ranks	14820 (87.2%)	2179 (12.8%)	16999
< 43 years old & rank < 7	13805 (94.4%)	812 (5.6%)	14617
< 43 years old & rank >=7	844 (42.2%)	1158 (57.8%)	2002
>= 43 years old	171 (45.0%)	209 (55.0%)	380

Note: Cancelled cycles are not included in the table.

Figure 1.3 Own and recipient fresh cycles: Evolution of total number of cycles



Note: Cancelled cycles are not included in the figure.

Section 2: own fresh cycles

Table 2.1 Own fresh cycles: Overview of cycles

Cycle	All Centres
Initiated	19061 (100.0%)
Cancelled	1715 (9.0%)
Aspiration	17346 (91.0%)
Embryo Transfer	15343 (80.5%)

Figure 2.2 Own fresh cycles: Female age and laborank

All Centres (N=16278, Missing=2783)




Figure 2.3 Own fresh cycles: Indications of ART




Table 2.4 Own fresh cycles: Indications of ART: female and male causes

	Statistic	All Centres
Female pathology	N	7970
Tubal	n/N (%)	3433/8408 (40.83%)
Endometriosis	n/N (%)	2316/8375 (27.65%)
Ovulatory	n/N (%)	1983/8151 (24.33%)
Other	n/N (%)	1865/8707 (21.42%)
Premature Ovarian Failure	n/N (%)	237/8436 (2.81%)
Genetic anomaly	n/N (%)	513/6260 (8.19%)
Immunological	n/N (%)	28/5930 (0.47%)
Male pathology	N	10644
Genetic anomaly	n/N (%)	471/9086 (5.18%)
Sperm abnormality	n/N (%)	11451/11731 (97.61%)
Immunological	n/N (%)	150/9692 (1.55%)

Some patients have more than one cause identified per cycle.

Figure 2.5 Own fresh cycles: Female age distribution




Figure 2.6 Own fresh cycles: Pituitary inhibition




Table 2.7 Own fresh cycles: Stimulation protocol

	Statistic	All Centres (N=19051, Missing=10)
Stimulation protocol		
Clomiphene	n/N (%)	1407/19051 (7.39%)
Gonadotrophins	n/N (%)	15909/19051 (83.51%)
Clomiphene + Gonadotrophins	n/N (%)	280/19051 (1.47%)
Aromatase Inhibitor + Gonadotrophins	n/N (%)	176/19051 (0.92%)
Substitution	n/N (%)	1/19051 (0.01%)
None	n/N (%)	659/19051 (3.46%)
Other	n/N (%)	619/19051 (3.25%)

Figure 2.8 Own fresh cycles: Total dose of Gonadotrophins (percentiles)





Figure 2.9 Own fresh cycles: Total dose of Gonadotrophins (boxplot)



Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.
 Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. + -sign indicates mean value.

Table 2.10 Own fresh cycles: Methods of fertilization

	Statistic	All Centres (N=16840, Missing=206)
Method of fertilization		
IVF	n/N (%)	4128/16840 (24.51%)
ICSI	n/N (%)	11771/16840 (69.90%)
Mixed (IVF + ICSI)	n/N (%)	941/16840 (5.59%)

1. Cycles with at least 1 oocyte retrieved are included.
2. Sperm of partner or donor are both included.

Table 2.11 Own fresh cycles: ICSI method sperm from partner

Sperm	All Centres (N=10669, Missing=467)					
	Fresh		Thawed		Total	
	N	%	N	%	N	%
Ejaculated	9401	93.91	610	6.09	10011	93.83
Surgical retrieved	235	35.71	423	64.29	658	6.17
Total	9636	90.32	1033	9.68	10669	100.00

Percentages are row percentages, except in the column 'Total'.

Table 2.12 Own fresh cycles: Transfers by age and rank categories

Age (yrs)	<36					[36-40[[40-43[Total	
	Rank	1	2	3-6	>=7	Total	1	2	3-6	>=7	Total	1	2	3-6	>=7		Total
All Centres (N=16278, Missing=1068)																	
Aspirations	4129	2409	2958	839	10335	1043	713	1223	592	3571	612	427	700	372	2111	261	16278
Transfers	3786	2243	2743	683	9455	940	646	1105	485	3176	529	382	622	286	1819	222	14672
Embryos transferred																	
1	3717	1326	671	258	5972	409	224	273	113	1019	152	100	170	54	476	61	7528
2	68	912	2050	333	3363	519	400	551	234	1704	223	149	185	92	649	63	5779
3	1	5	22	75	103	11	20	276	124	431	135	117	209	97	558	59	1151
>3	0	0	0	17	17	1	1	4	14	20	19	16	58	43	136	39	212
Unknown	0	0	0	0	0	0	1	1	0	2	0	0	0	0	0	0	2

Table 2.13 Own fresh cycles: Transfers by social security

All Centres (N=19061, Missing=0)			
	With social security	Without social security	Total
Initiated cycles	16211	2850	19061
Aspirations	14833	2513	17346
Transfers	13158	2185	15343
Embryos transferred			
1	7099	638	7737
2	5117	989	6106
3	839	408	1247
>3	101	149	250
Unknown	2	1	3

Figure 2.14 Own fresh cycles: Embryos transferred women < 36 years old

All Centres (N=9455, Missing=295)




Figure 2.15 Own fresh cycles: Embryos transferred women 36-40 years old

All Centres (N=3174, Missing=209)




Figure 2.16 Own fresh cycles: Embryos transferred women 40-43 years old

All Centres (N=1819, Missing=94)




Table 2.17 Own fresh cycles: Laboratory data

All Centres (N=19061, Missing=0)						
	Oocytes retrieved	Oocytes inseminated (IVF, ICSI or mixed)	2 PN oocytes	Transferred embryos	Cryopreserved embryos	
n	164354	141496	95690	24784	30448	
%	100.0%	86.1%	58.2%	15.1%	18.5%	
per initiated cycle	8.6	7.4	5.0	1.3	1.6	

Figure 2.18 Own fresh cycles: Summary pick-up cycles





Figure 2.19 Own fresh cycles: Distribution of embryo transfers

All Centres (N=15340, Missing=3)



Number of embryos transferred

1 embryo	: n (%) = 7737 (50.44%)
2 embryos	: n (%) = 6106 (39.80%)
3 embryos	: n (%) = 1247 (8.13%)
4 embryos	: n (%) = 185 (1.21%)
5 embryos	: n (%) = 39 (0.25%)
6 embryos	: n (%) = 24 (0.16%)
8 embryos	: n (%) = 1 (0.01%)


Table 2.20 Own fresh cycles: Cause of no transfer

	Statistic	All Centres
No Transfer	N	1950
No oocyte	n/N (%)	248/1950 (12.72%)
No sperm	n/N (%)	57/1950 (2.92%)
No transferable embryo available	n/N (%)	1221/1950 (62.62%)
OHSS risk	n/N (%)	107/1950 (5.49%)
Other reason	n/N (%)	344/1950 (17.64%)
Unknown	n/N (%)	80/1950 (4.10%)

Some patients can have more than one cause identified per cycle.

Figure 2.21 Own fresh cycles: Day of embryos transfer

All Centres (N=15318, Missing=25)



Day of Embryo Transfer






	Day 0: n (%) = 18 (0.12%)
	Day 1: n (%) = 34 (0.22%)
	Day 2: n (%) = 4071 (26.58%)
	Day 3: n (%) = 8842 (57.72%)
	Day 4-5-6-7: n (%) = 2353 (15.36%)

Table 2.22 Own fresh cycles: Cycles with cryopreservation

	All Centres (N=17020, Missing=26)
Number of cycles with cryopreservation	7226/17020 (42%)
Number of embryos cryopreserved	30448
Number of embryos per cryopreservation procedure	
Median	3.0
(Q1,Q3)	(2.0; 5.0)
Stage of the cryopreserved embryos	
2 PN	3623/30448 (12%)
Cleaved	22660/30448 (74%)
Blastocysts	4165/30448 (14%)
Percent freezing of non transferred embryos	30448/139570 (22%)

Based on all cycles with at least one oocyte retrieved.
Q1,Q3 = 1st and 3rd quartile.

Table 2.23 Own fresh cycles: Number of HCG+ pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
< 36 (yrs)					
All Centres (N=10335, Missing=435)					
Aspirations	4129	2409	2958	839	10335
Transfers	3786	2243	2743	683	9455
HCG + per aspiration cycle	1493/4119 (36.2%) (36.2% - 36.4%)	915/2399 (38.1%) (38.0% - 38.4%)	1051/2950 (35.6%) (35.5% - 35.8%)	277/837 (33.1%) (33.0% - 33.3%)	3736/10305 (36.3%) (36.1% - 36.4%)
HCG + per embryo transfer	1493/3776 (39.5%) (39.4% - 39.7%)	915/2233 (41.0%) (40.8% - 41.2%)	1051/2735 (38.4%) (38.3% - 38.6%)	277/681 (40.7%) (40.6% - 40.8%)	3736/9425 (39.6%) (39.5% - 39.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 2.23 Own fresh cycles: Number of HCG+ pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
[36-40[(yrs)					
All Centres (N=3571, Missing=330)					
Aspirations	1043	713	1223	592	3571
Transfers	940	646	1105	485	3176
HCG + per aspiration cycle	333/1038 (32.1%) (31.9% - 32.4%)	198/711 (27.8%) (27.8% - 28.1%)	331/1219 (27.2%) (27.1% - 27.4%)	193/590 (32.7%) (32.6% - 32.9%)	1055/3558 (29.7%) (29.5% - 29.9%)
HCG + per embryo transfer	333/935 (35.6%) (35.4% - 36.0%)	198/644 (30.7%) (30.7% - 31.0%)	331/1101 (30.1%) (30.0% - 30.3%)	193/483 (40.0%) (39.8% - 40.2%)	1055/3163 (33.4%) (33.2% - 33.6%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 2.23 Own fresh cycles: Number of HCG+ pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
[40-43] (yrs)					
All Centres (N=2111, Missing=199)					
Aspirations	612	427	700	372	2111
Transfers	529	382	622	286	1819
HCG + per aspiration cycle	140/610 (23.0%) (22.9% - 23.2%)	100/422 (23.7%) (23.4% - 24.6%)	139/699 (19.9%) (19.9% - 20.0%)	96/370 (25.9%) (25.8% - 26.3%)	475/2101 (22.6%) (22.5% - 23.0%)
HCG + per embryo transfer	140/527 (26.6%) (26.5% - 26.8%)	100/377 (26.5%) (26.2% - 27.5%)	139/621 (22.4%) (22.3% - 22.5%)	96/284 (33.8%) (33.6% - 34.3%)	475/1809 (26.3%) (26.1% - 26.7%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 2.23 Own fresh cycles: Number of HCG+ pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
>=43 (yrs)					
All Centres (N=261, Missing=101)					
Aspirations	48	35	40	138	261
Transfers	42	30	36	114	222
HCG + per aspiration cycle	2/45 (4.4%) (4.2% - 10.4%)	3/34 (8.8%) (8.6% - 11.4%)	2/37 (5.4%) (5.0% - 12.5%)	24/136 (17.6%) (17.4% - 18.8%)	31/252 (12.3%) (11.9% - 15.3%)
HCG + per embryo transfer	2/39 (5.1%) (4.8% - 11.9%)	3/29 (10.3%) (10.0% - 13.3%)	2/33 (6.1%) (5.6% - 13.9%)	24/112 (21.4%) (21.1% - 22.8%)	31/213 (14.6%) (14.0% - 18.0%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 2.24 Own fresh cycles: Number of clinical pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
< 36 (yrs)					
All Centres (N=10335, Missing=435)					
Aspirations	4129	2409	2958	839	10335
Transfers	3786	2243	2743	683	9455
Clinical Pregnancy per aspiration cycle	1300/4129 (31.5%) (31.5% - 31.5%)	780/2409 (32.4%) (32.4% - 32.4%)	909/2958 (30.7%) (30.7% - 30.7%)	204/839 (24.3%) (24.3% - 24.3%)	3193/10335 (30.9%) (30.9% - 30.9%)
Clinical Pregnancy per embryo transfer	1300/3786 (34.3%) (34.3% - 34.3%)	780/2243 (34.8%) (34.8% - 34.8%)	909/2743 (33.1%) (33.1% - 33.1%)	204/683 (29.9%) (29.9% - 29.9%)	3193/9455 (33.8%) (33.8% - 33.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.24 Own fresh cycles: Number of clinical pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
[36-40[(yrs)					
All Centres (N=3571, Missing=330)					
Aspirations	1043	713	1223	592	3571
Transfers	940	646	1105	485	3176
Clinical Pregnancy per aspiration cycle	286/1043 (27.4%) (27.4% - 27.4%)	160/713 (22.4%) (22.4% - 22.4%)	274/1223 (22.4%) (22.4% - 22.4%)	142/592 (24.0%) (24.0% - 24.0%)	862/3571 (24.1%) (24.1% - 24.1%)
Clinical Pregnancy per embryo transfer	286/940 (30.4%) (30.4% - 30.4%)	160/646 (24.8%) (24.8% - 24.8%)	274/1105 (24.8%) (24.8% - 24.8%)	142/485 (29.3%) (29.3% - 29.3%)	862/3176 (27.1%) (27.1% - 27.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.24 Own fresh cycles: Number of clinical pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
[40-43[(yrs)					
All Centres (N=2111, Missing=199)					
Aspirations	612	427	700	372	2111
Transfers	529	382	622	286	1819
Clinical Pregnancy per aspiration cycle	122/612 (19.9%) (19.9% - 19.9%)	87/427 (20.4%) (20.4% - 20.4%)	110/700 (15.7%) (15.7% - 15.7%)	64/372 (17.2%) (17.2% - 17.2%)	383/2111 (18.1%) (18.1% - 18.1%)
Clinical Pregnancy per embryo transfer	122/529 (23.1%) (23.1% - 23.1%)	87/382 (22.8%) (22.8% - 22.8%)	110/622 (17.7%) (17.7% - 17.7%)	64/286 (22.4%) (22.4% - 22.4%)	383/1819 (21.1%) (21.1% - 21.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.24 Own fresh cycles: Number of clinical pregnancies according to age and rank

Rank	1	2	3-6	>=7	Total
>=43 (yrs)					
All Centres (N=261, Missing=101)					
Aspirations	48	35	40	138	261
Transfers	42	30	36	114	222
Clinical Pregnancy per aspiration cycle	1/48 (2.1%) (2.1% - 2.1%)	3/35 (8.6%) (8.6% - 8.6%)	2/40 (5.0%) (5.0% - 5.0%)	14/138 (10.1%) (10.1% - 10.1%)	20/261 (7.7%) (7.7% - 7.7%)
Clinical Pregnancy per embryo transfer	1/42 (2.4%) (2.4% - 2.4%)	3/30 (10.0%) (10.0% - 10.0%)	2/36 (5.6%) (5.6% - 5.6%)	14/114 (12.3%) (12.3% - 12.3%)	20/222 (9.0%) (9.0% - 9.0%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.25 Own fresh cycles: Number of clinical pregnancies including FHB according to age and rank

Rank	1	2	3-6	>=7	Total
< 36 (yrs)					
All Centres (N=10335, Missing=435)					
Aspirations	4129	2409	2958	839	10335
Transfers	3786	2243	2743	683	9455
FHB: 1/2/3/4	1186/16/1/0	628/85/2/0	644/192/4/0	138/47/1/0	2596/340/8/0
Clinical Pregnancy + FHB per aspiration cycle	1203/4129 (29.1%) (29.1% - 29.1%)	715/2409 (29.7%) (29.7% - 29.7%)	840/2958 (28.4%) (28.4% - 28.4%)	186/839 (22.2%) (22.2% - 22.2%)	2944/10335 (28.5%) (28.5% - 28.5%)
Clinical Pregnancy + FHB per embryo transfer	1203/3786 (31.8%) (31.8% - 31.8%)	715/2243 (31.9%) (31.9% - 31.9%)	840/2743 (30.6%) (30.6% - 30.6%)	186/683 (27.2%) (27.2% - 27.2%)	2944/9455 (31.1%) (31.1% - 31.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.25 Own fresh cycles: Number of clinical pregnancies including FHB according to age and rank

Rank	1	2	3-6	>=7	Total
[36-40[(yrs)					
All Centres (N=3571, Missing=330)					
Aspirations	1043	713	1223	592	3571
Transfers	940	646	1105	485	3176
FHB: 1/2/3/4	215/36/1/0	117/21/2/0	202/40/2/1	98/19/4/0	632/116/9/1
Clinical Pregnancy + FHB per aspiration cycle	252/1043 (24.2%) (24.2% - 24.2%)	140/713 (19.6%) (19.6% - 19.6%)	245/1223 (20.0%) (20.0% - 20.0%)	121/592 (20.4%) (20.4% - 20.4%)	758/3571 (21.2%) (21.2% - 21.2%)
Clinical Pregnancy + FHB per embryo transfer	252/940 (26.8%) (26.8% - 26.8%)	140/646 (21.7%) (21.7% - 21.7%)	245/1105 (22.2%) (22.2% - 22.2%)	121/485 (24.9%) (24.9% - 24.9%)	758/3176 (23.9%) (23.9% - 23.9%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.25 Own fresh cycles: Number of clinical pregnancies including FHB according to age and rank

Rank	1	2	3-6	>=7	Total
[40-43[(yrs)					
All Centres (N=2111, Missing=199)					
Aspirations	612	427	700	372	2111
Transfers	529	382	622	286	1819
FHB: 1/2/3/4	98/7/2/0	59/13/1/0	82/7/1/0	44/10/0/0	283/37/4/0
Clinical Pregnancy + FHB per aspiration cycle	107/612 (17.5%) (17.5% - 17.5%)	73/427 (17.1%) (17.1% - 17.1%)	90/700 (12.9%) (12.9% - 12.9%)	54/372 (14.5%) (14.5% - 14.5%)	324/2111 (15.3%) (15.3% - 15.3%)
Clinical Pregnancy + FHB per embryo transfer	107/529 (20.2%) (20.2% - 20.2%)	73/382 (19.1%) (19.1% - 19.1%)	90/622 (14.5%) (14.5% - 14.5%)	54/286 (18.9%) (18.9% - 18.9%)	324/1819 (17.8%) (17.8% - 17.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.25 Own fresh cycles: Number of clinical pregnancies including FHB according to age and rank

Rank	1	2	3-6	>=7	Total
>=43 (yrs)					
All Centres (N=261, Missing=101)					
Aspirations	48	35	40	138	261
Transfers	42	30	36	114	222
FHB: 1/2/3/4	1/0/0/0	2/0/0/0	0/0/0/0	11/1/0/0	14/1/0/0
Clinical Pregnancy + FHB per aspiration cycle	1/48 (2.1%) (2.1% - 2.1%)	2/35 (5.7%) (5.7% - 5.7%)	0/40 (0.0% - 0.0%)	12/138 (8.7%) (8.7% - 8.7%)	15/261 (5.7%) (5.7% - 5.7%)
Clinical Pregnancy + FHB per embryo transfer	1/42 (2.4%) (2.4% - 2.4%)	2/30 (6.7%) (6.7% - 6.7%)	0/36 (0.0% - 0.0%)	12/114 (10.5%) (10.5% - 10.5%)	15/222 (6.8%) (6.8% - 6.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 2.26 Own fresh cycles: Number of deliveries according to age and rank

Rank	1	2	3-6	>=7	Total
< 36 (yrs)					
All Centres (N=10335, Missing=435)					
Aspirations	4129	2409	2958	839	10335
Transfers	3786	2243	2743	683	9455
Number per delivery: 1/2/3	956/12/0	514/70/0	543/143/3	87/23/0	2100/248/3
Delivery rate per aspiration cycle	968/4036 (24.0%) (23.4% - 25.7%)	584/2352 (24.8%) (24.2% - 26.6%)	689/2906 (23.7%) (23.3% - 25.1%)	110/733 (15.0%) (13.1% - 25.7%)	2351/10027 (23.4%) (22.7% - 25.7%)
Delivery rate per embryo transfer	968/3693 (26.2%) (25.6% - 28.0%)	584/2186 (26.7%) (26.0% - 28.6%)	689/2691 (25.6%) (25.1% - 27.0%)	110/577 (19.1%) (16.1% - 31.6%)	2351/9147 (25.7%) (24.9% - 28.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Table 2.26 Own fresh cycles: Number of deliveries according to age and rank

Rank	1	2	3-6	>=7	Total
[36-40[(yrs)					
All Centres (N=3571, Missing=330)					
Aspirations	1043	713	1223	592	3571
Transfers	940	646	1105	485	3176
Number per delivery: 1/2/3	170/25/1	96/16/1	161/30/1	73/12/1	500/83/4
Delivery rate per aspiration cycle	196/1029 (19.0%) (18.8% - 20.1%)	113/702 (16.1%) (15.8% - 17.4%)	192/1210 (15.9%) (15.7% - 16.8%)	86/542 (15.9%) (14.5% - 23.0%)	587/3483 (16.9%) (16.4% - 18.9%)
Delivery rate per embryo transfer	196/926 (21.2%) (20.9% - 22.3%)	113/635 (17.8%) (17.5% - 19.2%)	192/1092 (17.6%) (17.4% - 18.6%)	86/435 (19.8%) (17.7% - 28.0%)	587/3088 (19.0%) (18.5% - 21.3%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Table 2.26 Own fresh cycles: Number of deliveries according to age and rank

Rank	1	2	3-6	>=7	Total
[40-43[(yrs)					
All Centres (N=2111, Missing=199)					
Aspirations	612	427	700	372	2111
Transfers	529	382	622	286	1819
Number per delivery: 1/2/3	51/4/0	37/7/0	52/5/0	28/6/0	168/22/0
Delivery rate per aspiration cycle	55/603 (9.1%) (9.0% - 10.5%)	44/423 (10.4%) (10.3% - 11.2%)	57/695 (8.2%) (8.1% - 8.9%)	34/341 (10.0%) (9.1% - 17.5%)	190/2062 (9.2%) (9.0% - 11.3%)
Delivery rate per embryo transfer	55/520 (10.6%) (10.4% - 12.1%)	44/378 (11.6%) (11.5% - 12.6%)	57/617 (9.2%) (9.2% - 10.0%)	34/255 (13.3%) (11.9% - 22.7%)	190/1770 (10.7%) (10.4% - 13.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.


Table 2.26 Own fresh cycles: Number of deliveries according to age and rank

Rank	1	2	3-6	>=7	Total
>=43 (yrs)					
All Centres (N=261, Missing=101)					
Aspirations	48	35	40	138	261
Transfers	42	30	36	114	222
Number per delivery: 1/2/3	1/0/0	0/0/0	0/0/0	5/1/0	6/1/0
Delivery rate per aspiration cycle	1/48 (2.1%) (2.1% - 2.1%)	0/34 (0.0% - 2.9%)	0/40 (0.0% - 0.0%)	6/129 (4.7%) (4.3% - 10.9%)	7/251 (2.8%) (2.7% - 6.5%)
Delivery rate per embryo transfer	1/42 (2.4%) (2.4% - 2.4%)	0/29 (0.0% - 3.3%)	0/36 (0.0% - 0.0%)	6/105 (5.7%) (5.3% - 13.2%)	7/212 (3.3%) (3.2% - 7.7%)

NA=no cycles with data available.


In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Figure 2.27 Own fresh cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age




n/N (%) where n = Total number of uterine sacs; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.28 Own fresh cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age




n/N (%) where n = Total number of FHB; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.29 Own fresh cycles: Birth rate per transferred embryo according to age




n/N (%) where n = Total number of babies; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.30 Own fresh cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to rank




n/N (%) where n = Total number of uterine sacs; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.31 Own fresh cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to rank




n/N (%) where n = Total number of FHB; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.32 Own fresh cycles: Birth rate per transferred embryo according to rank




n/N (%) where n = Total number of babies; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 2.33 Own fresh cycles: Number of deliveries



Deliveries of twins or triplets are only counted once.

Figure 2.34 Own fresh cycles: Type of deliveries



Deliveries of twins or triplets are only counted once.

Table 2.35 Own fresh cycles: Sex of babies

All Centres (N=3593, Missing=24)	
Sex of baby	
Male	1738/3593 (48.37%)
Female	1732/3593 (48.20%)
Unknown	123/3593 (3.42%)

Table 2.36 Own fresh cycles: Birth weight


		All Centres	
		Statistic (N=3403, Missing=214)	
Birth weight (g)			
Singletons	N	2677	
	Mean	3251.8	
	Std	560.86	
	Median	3270.0	
	IQR	(2970.0; 3600.0)	
Twins	N	709	
	Mean	2368.6	
	Std	577.20	
	Median	2430.0	
	IQR	(2010.0; 2750.0)	
Triplets	N	17	
	Mean	1802.9	
	Std	595.23	
	Median	1715.0	
	IQR	(1330.0; 2000.0)	

Table 2.37 Own fresh cycles: Gestational age at delivery

	Statistic	All Centres (N=3193, Missing=38)
Gestational age at delivery (weeks)		
Singletons	N	2819
	Mean	39.0
	Std	2.10
	Median	39.4
	IQR	(38.3; 40.3)
Twins	N	367
	Mean	35.8
	Std	2.87
	Median	36.6
	IQR	(34.9; 37.7)
Triplets	N	7
	Mean	34.1
	Std	2.40
	Median	33.1
	IQR	(32.4; 37.6)

Twin or triplet birth is counted as one birth event.


Figure 2.38 Own fresh cycles: Birth weight (boxplot)



	Singleton	Twins	Triplets
N	2677	709	17
Missing	175	35	4
Mean	3251.8	2368.6	1802.9
SD	560.86	577.20	595.23
Median	3270.0	2430.0	1715.0
(Min,Max)	(550,5500)	(550,4550)	(1045,2995)
(Q1,Q3)	(2970,3600)	(2010,2750)	(1330,2000)

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.
 Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. +sign indicates mean value.

Figure 2.39 Own fresh cycles: Gestational age at delivery (boxplot)



	All Centres		
N	2819	367	7
Missing	33	5	0
Mean	39.0	35.8	34.1
SD	2.10	2.87	2.40
Median	39.4	36.6	33.1
(Min,Max)	(22,43)	(23,42)	(32,38)
(Q1,Q3)	(38,40)	(35,38)	(32,38)

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.

Q1, Q3 = 1st and 3rd quartile, IQR = Q3 - Q1. +sign indicates mean value.

Twin or triplet birth is counted as one birth event.

Table 2.40 Own fresh cycles: Prevalence of preterm birth according to type of delivery

Gestational age at delivery (weeks)	Type of delivery			
	Single birth event	Twin birth event	Triplet birth event	Total birth events
All Centres (N=3193, Missing=38)				
< 32	42 (1.5%)	34 (9.3%)	NA	76 (2.4%)
[32-37[207 (7.3%)	172 (46.9%)	5 (71.4%)	384 (12.0%)
>=37	2570 (91.2%)	161 (43.9%)	2 (28.6%)	2733 (85.6%)
Total	2819 (100.0%)	367 (100.0%)	7 (100.0%)	3193 (100.0%)

Twin or triplet birth is counted as one birth event.
 NA: no data available

Table 2.41 Own fresh cycles: Prevalence of low birth weight according to type of delivery

Birth weight (g)	Type of delivery				Total
	Singletons	Twins	Triplets		
All Centres (N=3403, Missing=214)					
< 1500	35 (1.3%)	53 (7.5%)	6 (35.3%)	94	(2.8%)
[1500-2500[158 (5.9%)	332 (46.8%)	8 (47.1%)	498	(14.6%)
>= 2500	2484 (92.8%)	324 (45.7%)	3 (17.6%)	2811	(82.6%)
Total	2677 (100.0%)	709 (100.0%)	17 (100.0%)	3403	(100.0%)

NA: no data available

Figure 2.42 Own fresh cycles: Evolution of number of embryos transferred





Figure 2.43 Own fresh cycles: Evolution of number of single and multiple deliveries



Section 3: Own cryo cycles

Table 3.1 Own cryo cycles: Overview of cryo cycles

Cryocycle	All Centres
Initiated	8576 (100.0%)
Cancelled	327 (3.8%)
Thawed	8249 (96.2%)
Embryo Transfer	6352 (74.1%)

Table 3.2 Own cryo cycles: Number of embryos transferred

	All Centres
Number of cycles with transfer	6352
Number of embryos transferred	
1	3065/6348 (48.28%)
2	3187/6348 (50.20%)
3	80/6348 (1.26%)
>3	16/6348 (0.25%)
Total number of embryos transferred	9745

Based on all cycles with at least one embryo transferred.

Table 3.3 Own cryo cycles: Pituitary inhibition

	Statistic	All Centres (N=8514, Missing=62)
Pituitary inhibition		
Yes	n/N (%)	470/8514 (5.52%)
No	n/N (%)	8044/8514 (94.48%)

Table 3.4 Own cryo cycles: Stimulation protocol

	Statistic	All Centres (N=8514, Missing=62)
Stimulation protocol		
Clomiphene	n/N (%)	945/8514 (11.10%)
Gonadotrophins	n/N (%)	297/8514 (3.49%)
Clomiphene + Gonadotrophins	n/N (%)	5/8514 (0.06%)
LHRH pump	n/N (%)	1/8514 (0.01%)
Substitution	n/N (%)	2137/8514 (25.10%)
None	n/N (%)	4585/8514 (53.85%)
Other	n/N (%)	544/8514 (6.39%)

Table 3.5 Own cryo cycles: Number of HCG+ pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=8576, Missing=0)					
Initiated cycles	6322	1590	532	132	8576
Thawed cycles	6099	1527	505	118	8249
Transfers	4712	1156	391	93	6352
HCG + per initiated cycle	1366/6299 (21.7%) (21.6% - 22.0%)	307/1585 (19.4%) (19.3% - 19.6%)	71/532 (13.3%) (13.3% - 13.3%)	20/129 (15.5%) (15.2% - 17.4%)	1764/8545 (20.6%) (20.6% - 20.9%)
HCG + per thawing cycle	1366/6076 (22.5%) (22.4% - 22.8%)	307/1522 (20.2%) (20.1% - 20.4%)	71/505 (14.1%) (14.1% - 14.1%)	20/115 (17.4%) (16.9% - 19.5%)	1764/8218 (21.5%) (21.4% - 21.8%)
HCG + per embryo transfer	1366/4695 (29.1%) (29.0% - 29.4%)	307/1152 (26.6%) (26.6% - 26.9%)	71/391 (18.2%) (18.2% - 18.2%)	20/90 (22.2%) (21.5% - 24.7%)	1764/6328 (27.9%) (27.8% - 28.1%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 3.6 Own cryo cycles: Number of clinical pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=8576, Missing=0)					
Initiated cycles	6322	1590	532	132	8576
Thawed cycles	6099	1527	505	118	8249
Transfers	4712	1156	391	93	6352
Clinical Pregnancy per initiated cycle	1160/6322 (18.3%) (18.3% - 18.3%)	259/1590 (16.3%) (16.3% - 16.3%)	62/532 (11.7%) (11.7% - 11.7%)	15/132 (11.4%) (11.4% - 11.4%)	1496/8576 (17.4%) (17.4% - 17.4%)
Clinical Pregnancy per thawing cycle	1160/6099 (19.0%) (19.0% - 19.0%)	259/1527 (17.0%) (17.0% - 17.0%)	62/505 (12.3%) (12.3% - 12.3%)	15/118 (12.7%) (12.7% - 12.7%)	1496/8249 (18.1%) (18.1% - 18.1%)
Clinical Pregnancy per embryo transfer	1160/4712 (24.6%) (24.6% - 24.6%)	259/1156 (22.4%) (22.4% - 22.4%)	62/391 (15.9%) (15.9% - 15.9%)	15/93 (16.1%) (16.1% - 16.1%)	1496/6352 (23.6%) (23.6% - 23.6%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 3.7 Own cryo cycles: Number of clinical pregnancies including FHB according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=8576, Missing=0)					
Initiated cycles	6322	1590	532	132	8576
Thawed cycles	6099	1527	505	118	8249
Transfers	4712	1156	391	93	6352
FHB: 1/2/3	867/149/5	210/22/0	48/5/0	12/1/0	1137/177/5
Clinical Pregnancy + FHB per initiated cycle	1021/6322 (16.1%) (16.1% - 16.1%)	232/1590 (14.6%) (14.6% - 14.6%)	53/532 (10.0%) (10.0% - 10.0%)	13/132 (9.8%) (9.8% - 9.8%)	1319/8576 (15.4%) (15.4% - 15.4%)
Clinical Pregnancy + FHB per thawing cycle	1021/6099 (16.7%) (16.7% - 16.7%)	232/1527 (15.2%) (15.2% - 15.2%)	53/505 (10.5%) (10.5% - 10.5%)	13/118 (11.0%) (11.0% - 11.0%)	1319/8249 (16.0%) (16.0% - 16.0%)
Clinical Pregnancy + FHB per embryo transfer	1021/4712 (21.7%) (21.7% - 21.7%)	232/1156 (20.1%) (20.1% - 20.1%)	53/391 (13.6%) (13.6% - 13.6%)	13/93 (14.0%) (14.0% - 14.0%)	1319/6352 (20.8%) (20.8% - 20.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.


Table 3.8 Own cryo cycles: Number of deliveries according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=8576, Missing=0)					
Initiated cycles	6322	1590	532	132	8576
Thawed cycles	6099	1527	505	118	8249
Transfers	4712	1156	391	93	6352
Number per delivery: 1/2/3	693/121/4	164/14/0	34/3/0	8/0/0	899/138/4
Delivery rate per initiated cycle	818/6248 (13.1%) (12.9% - 14.1%)	178/1567 (11.4%) (11.2% - 12.6%)	37/525 (7.0%) (7.0% - 8.3%)	8/129 (6.2%) (6.1% - 8.3%)	1041/8469 (12.3%) (12.1% - 13.4%)
Delivery rate per thawing cycle	818/6025 (13.6%) (13.4% - 14.6%)	178/1504 (11.8%) (11.7% - 13.2%)	37/498 (7.4%) (7.3% - 8.7%)	8/115 (7.0%) (6.8% - 9.3%)	1041/8142 (12.8%) (12.6% - 13.9%)
Delivery rate per embryo transfer	818/4644 (17.6%) (17.4% - 18.8%)	178/1134 (15.7%) (15.4% - 17.3%)	37/384 (9.6%) (9.5% - 11.3%)	8/90 (8.9%) (8.6% - 11.8%)	1041/6252 (16.7%) (16.4% - 18.0%)

NA=no cycles with data available.


In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Figure 3.9 Own cryo cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age




n/N (%) where n = Total number of uterine sacs; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 3.10 Own cryo cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age




n/N (%) where n = Total number of FHB; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 3.11 Own cryo cycles: Birth rate per transferred embryo according to age




n/N (%) where n = Total number of babies; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 3.12 Own cryo cycles: Number of deliveries



Deliveries of twins or triplets are only counted once.

Figure 3.13 Own cryo cycles: Type of deliveries



Deliveries of twins or triplets are only counted once.

Table 3.14 Own cryo cycles: Sex of babies

All Centres (N=1184, Missing=3)	
Sex of baby	
Male	575/1184 (48.56%)
Female	574/1184 (48.48%)
Unknown	35/1184 (2.96%)

Table 3.15 Own cryo cycles: Birth weight


	Statistic	All Centres (N=1113, Missing=74)
Birth weight (g)		
Singletons	N	840
	Mean	3364.2
	Std	577.79
	Median	3390.0
	IQR	(3100.0; 3700.0)
Twins	N	262
	Mean	2553.4
	Std	514.52
	Median	2560.0
	IQR	(2225.0; 2850.0)
Triplets	N	11
	Mean	1891.8
	Std	535.77
	Median	2040.0
	IQR	(1200.0; 2290.0)

Table 3.16 Own cryo cycles: Gestational age at delivery

	Statistic	All Centres (N=1034, Missing=7)
Gestational age at delivery (weeks)		
Singletons	N	892
	Mean	39.1
	Std	2.47
	Median	39.4
	IQR	(38.4; 40.4)
Twins	N	138
	Mean	36.2
	Std	2.29
	Median	36.9
	IQR	(35.0; 37.9)
Triplets	N	4
	Mean	33.5
	Std	2.87
	Median	33.7
	IQR	(31.6; 35.3)


Twin or triplet birth is counted as one birth event.

Figure 3.17 Own cryo cycles: Birth weight (boxplot)



Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.
 $Q1, Q3 = 1st \text{ and } 3rd \text{ quartile, } IQR = Q3 - Q1. +\text{-sign indicates mean value.}$

Figure 3.18 Own cryo cycles: Gestational age at delivery (boxplot)



	All Centres		
N	892	138	4
Missing	7	0	0
Mean	39.1	36.2	33.5
SD	2.47	2.29	2.87
Median	39.4	36.9	33.7
(Min,Max)	(2,42)	(26,41)	(30,37)
(Q1,Q3)	(38,40)	(35,38)	(32,35)

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.

Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. +sign indicates mean value.

Twin or triplet birth is counted as one birth event.

Table 3.19 Own cryo cycles: Prevalence of preterm birth according to type of delivery

Gestational age at delivery (weeks)	Type of delivery			
	Single birth event	Twin birth event	Triplet birth event	Total birth events
All Centres (N=1034, Missing=7)				
< 32	15 (1.7%)	5 (3.6%)	1 (25.0%)	21 (2.0%)
[32-37[60 (6.7%)	65 (47.1%)	3 (75.0%)	128 (12.4%)
>=37	817 (91.6%)	68 (49.3%)	NA	885 (85.6%)
Total	892 (100.0%)	138 (100.0%)	4 (100.0%)	1034 (100.0%)

Twin or triplet birth is counted as one birth event.
 NA: no data available

Table 3.20 Own cryo cycles: Prevalence of low birth weight according to type of delivery

Birth weight (g)	Type of delivery				Total
	Singletons	Twins	Triplets		
All Centres (N=1113, Missing=74)					
< 1500	10 (1.2%)	6 (2.3%)	3 (27.3%)	19	(1.7%)
[1500-2500[36 (4.3%)	111 (42.4%)	7 (63.6%)	154	(13.8%)
>= 2500	794 (94.5%)	145 (55.3%)	1 (9.1%)	940	(84.5%)
Total	840 (100.0%)	262 (100.0%)	11 (100.0%)	1113	(100.0%)

NA: no available data

Section 4: Fresh recipient cycles

Table 4.1 Fresh recipient cycles: Overview of cycles

Cycle	All Centres
Initiated	820 (100.0%)
Cancelled	89 (10.9%)
At least one oocyte received	731 (89.1%)
Embryo Transfer	645 (78.7%)

Figure 4.2 Fresh recipient cycles: Female age and laborank

All Centres (N=650, Missing=170)




Figure 4.3 Fresh recipient cycles: Female age distribution





Figure 4.4 Fresh recipient cycles: Pituitary inhibition

All Centres (N=820, Missing=0)



Pituitary Inhibition





	Agonist - long : n (%) = 364 (44.39%)
	Agonist - short : n (%) = 8 (0.98%)
	Antagonist : n (%) = 4 (0.49%)
	None : n (%) = 444 (54.15%)

Table 4.5 Fresh recipient cycles: Stimulation protocol

	Statistic	All Centres (N=820, Missing=0)
Stimulation protocol		
Gonadotrophins	n/N (%)	2/820 (0.24%)
Substitution	n/N (%)	706/820 (86.10%)
None	n/N (%)	27/820 (3.29%)
Other	n/N (%)	85/820 (10.37%)

Table 4.6 Fresh recipient cycles: Number of HCG+ pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=820, Missing=0)					
Initiated cycles	300	186	185	149	820
At least one oocyte received	274	168	159	130	731
Transfers	236	153	135	121	645
HCG + per initiated cycle	79/299 (26.4%) (26.3% - 26.7%)	54/184 (29.3%) (29.0% - 30.1%)	40/181 (22.1%) (21.6% - 23.8%)	51/147 (34.7%) (34.2% - 35.6%)	224/811 (27.6%) (27.3% - 28.4%)
HCG + per cycles with at least one oocyte received	79/273 (28.9%) (28.8% - 29.2%)	54/166 (32.5%) (32.1% - 33.3%)	40/155 (25.8%) (25.2% - 27.7%)	51/128 (39.8%) (39.2% - 40.8%)	224/722 (31.0%) (30.6% - 31.9%)
HCG + per embryo transfer	79/236 (33.5%) (33.5% - 33.5%)	54/151 (35.8%) (35.3% - 36.6%)	40/132 (30.3%) (29.6% - 31.9%)	51/120 (42.5%) (42.1% - 43.0%)	224/639 (35.1%) (34.7% - 35.7%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 4.7 Fresh recipient cycles: Number of clinical pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=820, Missing=0)					
Initiated cycles	300	186	185	149	820
At least one oocyte received	274	168	159	130	731
Transfers	236	153	135	121	645
Clinical Pregnancy per initiated cycle	62/300 (20.7%) (20.7% - 20.7%)	45/186 (24.2%) (24.2% - 24.2%)	32/185 (17.3%) (17.3% - 17.3%)	40/149 (26.8%) (26.8% - 26.8%)	179/820 (21.8%) (21.8% - 21.8%)
Clinical Pregnancy per cycles with at least one oocyte received	62/274 (22.6%) (22.6% - 22.6%)	45/168 (26.8%) (26.8% - 26.8%)	32/159 (20.1%) (20.1% - 20.1%)	40/130 (30.8%) (30.8% - 30.8%)	179/731 (24.5%) (24.5% - 24.5%)
Clinical Pregnancy per embryo transfer	62/236 (26.3%) (26.3% - 26.3%)	45/153 (29.4%) (29.4% - 29.4%)	32/135 (23.7%) (23.7% - 23.7%)	40/121 (33.1%) (33.1% - 33.1%)	179/645 (27.8%) (27.8% - 27.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 4.8 Fresh recipient cycles: Number of clinical pregnancies including FHB according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=820, Missing=0)					
Initiated cycles	300	186	185	149	820
At least one oocyte received	274	168	159	130	731
Transfers	236	153	135	121	645
FHB: 1/2/3	43/11/0	34/6/1	20/7/1	23/13/1	120/37/3
Clinical Pregnancy + FHB per initiated cycle	54/300 (18.0%) (18.0% - 18.0%)	41/186 (22.0%) (22.0% - 22.0%)	28/185 (15.1%) (15.1% - 15.1%)	37/149 (24.8%) (24.8% - 24.8%)	160/820 (19.5%) (19.5% - 19.5%)
Clinical Pregnancy + FHB per cycles with at least one oocyte received	54/274 (19.7%) (19.7% - 19.7%)	41/168 (24.4%) (24.4% - 24.4%)	28/159 (17.6%) (17.6% - 17.6%)	37/130 (28.5%) (28.5% - 28.5%)	160/731 (21.9%) (21.9% - 21.9%)
Clinical Pregnancy + FHB per embryo transfer	54/236 (22.9%) (22.9% - 22.9%)	41/153 (26.8%) (26.8% - 26.8%)	28/135 (20.7%) (20.7% - 20.7%)	37/121 (30.6%) (30.6% - 30.6%)	160/645 (24.8%) (24.8% - 24.8%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.


Table 4.9 Fresh recipient cycles: Number of deliveries according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=820, Missing=0)					
Initiated cycles	300	186	185	149	820
At least one oocyte received	274	168	159	130	731
Transfers	236	153	135	121	645
Number per delivery: 1/2/3	29/9/0	26/3/0	16/5/0	17/6/0	88/23/0
Delivery rate per initiated cycle	38/294 (12.9%) (12.7% - 14.7%)	29/178 (16.3%) (15.6% - 19.9%)	21/177 (11.9%) (11.4% - 15.7%)	23/134 (17.2%) (15.4% - 25.5%)	111/783 (14.2%) (13.5% - 18.0%)
Delivery rate per cycles with at least one oocyte received	38/268 (14.2%) (13.9% - 16.1%)	29/160 (18.1%) (17.3% - 22.0%)	21/151 (13.9%) (13.2% - 18.2%)	23/115 (20.0%) (17.7% - 29.2%)	111/694 (16.0%) (15.2% - 20.2%)
Delivery rate per embryo transfer	38/231 (16.5%) (16.1% - 18.2%)	29/145 (20.0%) (19.0% - 24.2%)	21/128 (16.4%) (15.6% - 20.7%)	23/107 (21.5%) (19.0% - 30.6%)	111/611 (18.2%) (17.2% - 22.5%)

NA=no cycles with data available.


In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Figure 4.10 Fresh recipient cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age




n/N (%) where n = Total number of uterine sacs; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 4.11 Fresh recipient cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age




n/N (%) where n = Total number of FHB; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 4.12 Fresh recipient cycles: Birth rate per transferred embryo according to age



n/N (%) where n = Total number of babies; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.


Figure 4.13 Fresh recipient cycles: Number of deliveries



Deliveries of twins or triplets are only counted once.

Table 4.14 Fresh recipient cycles: Type of deliveries

All Centres (N=111, Missing=0)



Deliveries of twins or triplets are only counted once.

Table 4.15 Fresh recipient cycles: Sex of babies

All Centres (N=133, Missing=1)	
Sex of baby	
Male	65/133 (48.87%)
Female	55/133 (41.35%)
Unknown	13/133 (9.77%)

Table 4.16 Fresh recipient cycles: Birth weight


	Statistic	All Centres (N=118, Missing=16)
Birth Weight (g)		
Singletons	N	73
	Mean	3278.0
	Std	708.87
	Median	3380.0
	IQR	(3040.0; 3720.0)
Twins	N	45
	Mean	2328.8
	Std	486.26
	Median	2430.0
	IQR	(2010.0; 2660.0)

Table 4.17 Fresh recipient cycles: Gestational age at delivery

	Statistic	All Centres (N=110, Missing=1)
Gestational age at delivery (weeks)		
Singletons	N	87
	Mean	38.2
	Std	2.72
	Median	38.7
	IQR	(37.3; 40.1)
Twins	N	23
	Mean	35.5
	Std	2.72
	Median	36.4
	IQR	(34.1; 37.7)

Twin or triplet birth is counted as one birth event.


Figure 4.18 Fresh recipient cycles: Birth weight (boxplot)



	Singleton	Twins	Triplets
N	73	45	
Missing	15	1	
Mean	3278.0	2328.8	
SD	708.87	486.26	
Median	3380.0	2430.0	
(Min,Max)	(761,4520)	(900,3080)	
(Q1,Q3)	(3040,3720)	(2010,2660)	

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.
 Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. +-sign indicates mean value.

Figure 4.19 Fresh recipient cycles: Gestational age at delivery (boxplot)



	Singleton	Twins	Triplets
All Centres			
N	87	23	
Missing	1	0	
Mean	38.2	35.5	
SD	2.72	2.72	
Median	38.7	36.4	
(Min,Max)	(26,42)	(27,39)	
(Q1,Q3)	(37,40)	(34,38)	

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.

Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. +-sign indicates mean value.

Twin or triplet birth is counted as one birth event.

Table 4.20 Fresh recipient cycles: Prevalence of preterm birth according to type of delivery

Gestational age at delivery (weeks)	Type of delivery			Total birth events
	Single birth event	Twin birth event	Triplet birth event	
All Centres (N=110, Missing=1)				
< 32	2 (2.3%)	1 (4.3%)	NA	3 (2.7%)
[32-37[16 (18.4%)	13 (56.5%)	NA	29 (26.4%)
>=37	69 (79.3%)	9 (39.1%)	NA	78 (70.9%)
Total	87 (100.0%)	23 (100.0%)	NA	110 (100.0%)

Twin or triplet birth is counted as one birth event.
 NA: no data available

Table 4.21 Fresh recipient cycles: Prevalence of low birth weight according to type of delivery

Birth weight (g)	Type of delivery			Total
	Singletons	Twins	Triplets	
All Centres (N=118, Missing=16)				
< 1500	3 (4.1%)	3 (6.7%)	NA	6 (5.1%)
[1500-2500[5 (6.8%)	24 (53.3%)	NA	29 (24.6%)
>= 2500	65 (89.0%)	18 (40.0%)	NA	83 (70.3%)
Total	73 (100.0%)	45 (100.0%)	NA	118 (100.0%)

NA: no data available

Section 5: Cryo recipient cycles

Table 5.1 Cryo recipient cycles: Overview of cryo cycles

Cryo cycle	All Centres	
Initiated	294	(100.0%)
Cancelled	3	(1.0%)
Thawed	291	(99.0%)
Embryo Transfer	231	(78.6%)

Table 5.2 Cryo recipient cycles: Number of embryos transferred

	All Centres
Number of cycles with transfer	231
Number of embryos transferred	
1	91/229 (39.74%)
2	121/229 (52.84%)
3	14/229 (6.11%)
>3	3/229 (1.31%)
Total number of embryos transferred	387

Based on all cycles with at least one embryo transferred.

Table 5.3 Cryo recipient cycles: Pituitary inhibition

	Statistic	All Centres (N=294, Missing=0)
Pituitary inhibition		
Yes	n/N (%)	19/294 (6.46%)
No	n/N (%)	275/294 (93.54%)

Table 5.4 Cryo recipient cycles: Stimulation protocol

	Statistic	All Centres (N=294, Missing=0)
Stimulation protocol		
Clomiphene	n/N (%)	14/294 (4.76%)
Gonadotrophins	n/N (%)	2/294 (0.68%)
Substitution	n/N (%)	194/294 (65.99%)
None	n/N (%)	43/294 (14.63%)
Other	n/N (%)	41/294 (13.95%)

Table 5.5 Cryo recipient cycles: Number of HCG+ pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=294, Missing=0)					
Initiated cycles	129	67	48	50	294
Thawing cycles	129	66	46	50	291
Transfers	97	49	39	46	231
HCG + per initiated cycle	23/127 (18.1%) (17.8% - 19.4%)	13/64 (20.3%) (19.4% - 23.9%)	11/47 (23.4%) (22.9% - 25.0%)	13/49 (26.5%) (26.0% - 28.0%)	60/287 (20.9%) (20.4% - 22.8%)
HCG + per thawing cycles	23/127 (18.1%) (17.8% - 19.4%)	13/63 (20.6%) (19.7% - 24.2%)	11/45 (24.4%) (23.9% - 26.1%)	13/49 (26.5%) (26.0% - 28.0%)	60/284 (21.1%) (20.6% - 23.0%)
HCG + per embryo transfer	23/95 (24.2%) (23.7% - 25.8%)	13/46 (28.3%) (26.5% - 32.7%)	11/38 (28.9%) (28.2% - 30.8%)	13/45 (28.9%) (28.3% - 30.4%)	60/224 (26.8%) (26.0% - 29.0%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing HCG results as negative and positive, respectively.

Table 5.6 Cryo recipient cycles: Number of clinical pregnancies according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=294, Missing=0)					
Initiated cycles	129	67	48	50	294
Thawing cycles	129	66	46	50	291
Transfers	97	49	39	46	231
Clinical Pregnancy per initiated cycle	20/129 (15.5%) (15.5% - 15.5%)	11/67 (16.4%) (16.4% - 16.4%)	10/48 (20.8%) (20.8% - 20.8%)	11/50 (22.0%) (22.0% - 22.0%)	52/294 (17.7%) (17.7% - 17.7%)
Clinical Pregnancy per thawing cycles	20/129 (15.5%) (15.5% - 15.5%)	11/66 (16.7%) (16.7% - 16.7%)	10/46 (21.7%) (21.7% - 21.7%)	11/50 (22.0%) (22.0% - 22.0%)	52/291 (17.9%) (17.9% - 17.9%)
Clinical Pregnancy per embryo transfer	20/97 (20.6%) (20.6% - 20.6%)	11/49 (22.4%) (22.4% - 22.4%)	10/39 (25.6%) (25.6% - 25.6%)	11/46 (23.9%) (23.9% - 23.9%)	52/231 (22.5%) (22.5% - 22.5%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.

Table 5.7 Cryo recipient cycles: Number of clinical pregnancies including FHB according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=294, Missing=0)					
Initiated cycles	129	67	48	50	294
Thawing cycles	129	66	46	50	291
Transfers	97	49	39	46	231
FHB: 1/2/3	15/1	8/2	5/3	7/2	35/8
Clinical Pregnancy + FHB per initiated cycle	16/129 (12.4%) (12.4% - 12.4%)	10/67 (14.9%) (14.9% - 14.9%)	8/48 (16.7%) (16.7% - 16.7%)	9/50 (18.0%) (18.0% - 18.0%)	43/294 (14.6%) (14.6% - 14.6%)
Clinical Pregnancy + FHB per thawing cycles	16/129 (12.4%) (12.4% - 12.4%)	10/66 (15.2%) (15.2% - 15.2%)	8/46 (17.4%) (17.4% - 17.4%)	9/50 (18.0%) (18.0% - 18.0%)	43/291 (14.8%) (14.8% - 14.8%)
Clinical Pregnancy + FHB per embryo transfer	16/97 (16.5%) (16.5% - 16.5%)	10/49 (20.4%) (20.4% - 20.4%)	8/39 (20.5%) (20.5% - 20.5%)	9/46 (19.6%) (19.6% - 19.6%)	43/231 (18.6%) (18.6% - 18.6%)

NA=no cycles with data available.

In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing results as negative and positive, respectively.


Table 5.8 Cryo recipient cycles: Number of deliveries according to age

Age (yrs.)	< 36	[36-40[[40-43[>=43	All ages
All Centres (N=294, Missing=0)					
Initiated cycles	129	67	48	50	294
Thawing cycles	129	66	46	50	291
Transfers	97	49	39	46	231
Number per delivery: 1/2/3	11/1/0	7/0/0	2/1/0	5/2/0	25/4/0
Delivery rate per initiated cycle	12/127 (9.4%) (9.3% - 10.9%)	7/66 (10.6%) (10.4% - 11.9%)	3/48 (6.3%) (6.3% - 6.3%)	7/50 (14.0%) (14.0% - 14.0%)	29/291 (10.0%) (9.9% - 10.9%)
Delivery rate per thawing cycles	12/127 (9.4%) (9.3% - 10.9%)	7/65 (10.8%) (10.6% - 12.1%)	3/46 (6.5%) (6.5% - 6.5%)	7/50 (14.0%) (14.0% - 14.0%)	29/288 (10.1%) (10.0% - 11.0%)
Delivery rate per embryo transfer	12/95 (12.6%) (12.4% - 14.4%)	7/48 (14.6%) (14.3% - 16.3%)	3/39 (7.7%) (7.7% - 7.7%)	7/46 (15.2%) (15.2% - 15.2%)	29/228 (12.7%) (12.6% - 13.9%)

NA=no cycles with data available.


In the calculation of the ratios, only cycles with available data are considered. In the line underneath, the range expresses the minimum and maximum possible rates when accounting for missing data by considering missing delivery as negative and positive, respectively.

Figure 5.9 Cryo recipient cycles: Implantation rate (No. of uterine sacs) per transferred embryo according to age




n/N (%) where n = Total number of uterine sacs; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 5.10 Cryo recipient cycles: Clinical implantation rate (No. of FHB) per transferred embryo according to age




n/N (%) where n = Total number of FHB; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.

Figure 5.11 Cryo recipient cycles: Birth rate per transferred embryo according to age



n/N (%) where n = Total number of babies; N = Total number of embryos transferred; %= n*100/N; NA = No cycles with data available.


Figure 5.12 Cryo recipient cycles: Number of deliveries



Deliveries of twins or triplets are only counted once.

Table 5.13 Cryo recipient cycles: Type of deliveries

All Centres (N=28, Missing=1)



Deliveries of twins or triplets are only counted once.

Table 5.14 Cryo recipient cycles: Sex of babies

All Centres (N=32, Missing=1)	
Sex of baby	
Male	10/32 (31.25%)
Female	18/32 (56.25%)
Unknown	4/32 (12.50%)

Table 5.15 Cryo recipient cycles: Birth weight


	Statistic	All Centres (N=27, Missing=6)
Birth Weight (g)		
Singletons	N	19
	Mean	3201.1
	Std	548.48
	Median	3010.0
	IQR	(2830.0; 3480.0)
Twins	N	8
	Mean	2154.4
	Std	597.41
	Median	2017.5
	IQR	(1720.0; 2775.0)

Table 5.16 Cryo recipient cycles: Gestational age at delivery

	Statistic	All Centres (N=26, Missing=3)
Gestational age at delivery (weeks)		
Singletons	N	22
	Mean	38.9
	Std	1.78
	Median	39.1
	IQR	(37.9; 40.1)
Twins	N	4
	Mean	35.1
	Std	1.77
	Median	35.4
	IQR	(33.6; 36.6)

Twin or triplet birth is counted as one birth event.


Figure 5.17 Cryo recipient cycles: Birth weight (boxplot)



	Singleton	Twins	Triplets
N	19	8	
Missing	6	0	
Mean	3201.1	2154.4	
SD	548.48	597.41	
Median	3010.0	2017.5	
(Min,Max)	(2330,4460)	(1310,2900)	
(Q1,Q3)	(2830,3480)	(1720,2775)	

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.
 Q1, Q3 = 1st and 3rd quartile, IQR = $Q3 - Q1$. +sign indicates mean value.

Figure 5.18 Cryo recipient cycles: Gestational age at delivery (boxplot)



	Singleton	Twins	Triplets
All Centres			
N	22	4	
Missing	3	0	
Mean	38.9	35.1	
SD	1.78	1.77	
Median	39.1	35.4	
(Min,Max)	(35,42)	(33,37)	
(Q1,Q3)	(38,40)	(34,37)	

Box plot shows median and interquartile range. Whiskers are drawn at $(Q3+1.5*IQR, Q1-1.5*IQR)$.

Q1, Q3 = 1st and 3rd quartile, $IQR = Q3 - Q1$. +-sign indicates mean value.

Twin or triplet birth is counted as one birth event.

Table 5.19 Cryo recipient cycles: Prevalence of preterm birth according to type of delivery

Gestational age at delivery (weeks)	Type of delivery			Total birth events
	Single birth event	Twin birth event	Triplet birth event	
All Centres (N=26, Missing=3)				
< 32	NA	NA	NA	NA
[32-37[4 (18.2%)	4 (100.0%)	NA	8 (30.8%)
>=37	18 (81.8%)	NA	NA	18 (69.2%)
Total	22 (100.0%)	4 (100.0%)	NA	26 (100.0%)

Twin or triplet birth is counted as one birth event.

Table 5.20 Cryo recipient cycles: Prevalence of low birth weight according to type of delivery

Birth weight (g)	Type of delivery			Total
	Singletons	Twins	Triplets	
All Centres (N=27, Missing=6)				
< 1500	NA	1 (12.5%)	NA	1 (3.7%)
[1500-2500[1 (5.3%)	4 (50.0%)	NA	5 (18.5%)
>= 2500	18 (94.7%)	3 (37.5%)	NA	21 (77.8%)
Total	19 (100.0%)	8 (100.0%)	NA	27 (100.0%)

NA: no data available

Section 6: Fresh donor cycles

Table 6.1 Fresh donor cycles: Overview of cycles

Cycle	All Centres
Initiated	578 (100.0%)
Cancelled	45 (7.8%)
At least one oocyte received	533 (92.2%)

Figure 6.2 Fresh donor cycles: Female age distribution




Table 6.3 Fresh donor cycles: Pituitary inhibition





Table 6.4 Fresh donor cycles: Stimulation protocol

	Statistic	All Centres (N=578, Missing=0)
Stimulation protocol		
Gonadotrophins	n/N (%)	542/578 (93.77%)
Clomiphene + Gonadotrophins	n/N (%)	1/578 (0.17%)
Substitution	n/N (%)	1/578 (0.17%)
None	n/N (%)	18/578 (3.11%)
Other	n/N (%)	16/578 (2.77%)

Figure 6.5 Fresh donor cycles: Total dose of Gonadotrophins (percentiles)



Section 7: Appendix

Table 7.1 : Definitions

Term	Definition
Clinical pregnancy	The presence of intra- or extra-uterine sacs on an ultrasound scan.
Delivery	Birth of a child, death or alive, of $\geq 500\text{g}$ or ≥ 22 weeks if birth weight is unknown.
Gestational age	Age of an embryo or fetus calculated by adding 14 days (2 weeks) to the number of completed weeks since fertilization.

Table 7.2 : List of B-centres having supplied data

City	Centre
Antwerpen	Dienst Fertiliteit, Algemeen Ziekenhuis Middelheim
Braine L'alleud	Centre de Fécondation ,C.H. Interrégional Edith Cavell (CHIREC)
Brugge	BIRTH - Fertilitetskliniek, Algemeen Ziekenhuis Sint-Jan
Brussel	Centrum voor Reproductieve Geneeskunde, UZ-Brussel
Bruxelles	Clinique de Procréation Médicalement Assistée, Hôpital Universitaire Saint- Pierre – U.L.B.
Bruxelles	Service de Gynécologie, Cliniques Universitaires Saint-Luc – U.C.L.
Bruxelles	Centre de FIV de l'ULB- Hôpital Erasme
Charleroi	Service Gyn/Obst,Clinique Notre Dame
Edegem	Centrum voor Reproductieve Geneeskunde, Universitair Ziekenhuis Antwerpen - U.I.A.
Genk	Centre for Reproductive Medicine, Ziekenhuis Oost-Limburg - St. Jan
Gent	Vrouwenkliniek - Infertiliteitscentrum, U.Z. – Gent
Gent	Centrum voor Fertilitetstherapie, A.Z. Jan Palfijn
Leuven	Dienst Gynaecologie, Universitaire Ziekenhuizen K.U.Leuven Gasthuisberg
Leuven	Unit Reproductieve Geneeskunde, Regionaal Ziekenhuis Heilig Hart
Libramont	Centre d'Infertilité, Centre Hospitalier de l'Ardenne
Liège	Centre de FIV, Centre Hospitalier Régional de la Citadelle
Namur	Service Gynéco, Centre Hospitalier Régional de Namur
Rocourt	Centre Liégeois pour l'Etude et le Traitement de la Stérilité, Clinique Saint Vincent

Colophon

College van Geneesheren "Reproductieve Geneeskunde"/

Collège de Médecins "Médecine de la Reproduction"

B. Lejeune, President

M. Camus, Vice-President

F. Vandekerckhove, Secretary

C. Wyns, Secretary

A. Delbaere, Member

T. D'Hooghe, Member

M. Dubois, Member

W. Ombelet, Member

Data handling and analysis

Interuniversity Institute for Biostatistics and statistical Bioinformatics

Katholieke Universiteit Leuven & Universiteit Hasselt

A. Belmans, K. Bogaerts, E. Lesaffre

Ecole de Santé Publique

Université de Liège

A. Albert, N. Gillain, E. Husson

This report is electronically available at www.belrap.be.